

**AMINOALKYL-SUBSTITUTED AROMATIC BICYCLIC COMPOUNDS,
METHODS FOR THEIR PREPARATION AND THEIR USE AS
PHARMACEUTICALS**

5 This application claims priority to German Patent Application 10139416.0, filed August 17, 2001, which is hereby incorporated by reference, in its entirety. All references cited below, including patents, patent applications and scientific journals and books also are herein incorporated by reference in their entirety.

10 **FIELD OF THE INVENTION**

The invention relates to aminoalkyl-substituted aromatic bicyclic compounds and to the physiologically acceptable salts and physiologically functional derivatives thereof.


15 **BACKGROUND OF THE INVENTION**

Structurally similar nonaromatic bicyclic compounds with pharmacological action have already been described in the prior art (for example in WO 01/21577).

20 The present invention provides compounds which cause a reduction in weight in mammals and which are suitable for preventing and treating obesity and diabetes.

SUMMARY OF THE INVENTION


25 The present invention relates to aminoalkyl-substituted aromatic bicyclic compounds of formula I,



wherein A, X, D, E, G, L, B, R5, R1, R2, R3, W, U, T, Y, R6 and R7 have the meanings as indicated herein. The compounds of formula I are valuable pharmaceutically active compounds which are suitable, for example, for the treatment of obesity, type II diabetes, arteriosclerosis, high blood pressure, paresthesia, depression, anxiety, anxiety neuroses, schizophrenia, disorders associated with the circadian rhythm, and drug abuse, as well as normalizing lipid metabolism.

DETAILED DESCRIPTION OF THE EMBODIMENTS

The invention therefore relates to compounds of formula I,



I

in which

A is (C₁-C₈)alkyl, (C₀-C₈)alkylenearyl, or a 3- to 12-membered mono- or bicyclic ring which may contain one or more heteroatoms selected from the group consisting of N, O and S and the 3- to 12-membered ring may carry further substituents, such as F, Cl, Br, NO₂, CF₃, OCF₃, CN, (C₁-C₆)alkyl, aryl, CON(R₃₇)(R₃₈), N(R₃₉)(R₄₀), OH, O-(C₁-C₆)alkyl, S-(C₁-C₆)alkyl, or NHCO(C₁-C₆)alkyl;

X is a bond, C(R₈)(R₉), C(OR₁₀)(R₁₁), O, N(R₁₂), S, SO, SO₂, or CO; wherein R₈, R₉, R₁₀, R₁₁, R₁₂ are, independently of one another, H, (C₁-C₆)alkyl;

D is N, or C(R₄₁);

E is N, or C(R42);

G is N, or C(R43);

5 L is N, or C(R44);

R1, R2, R3, R41, R42, R43, R44 are, independently of one another, H, F, Cl, Br, J, OH, CF₃, NO₂, CN, OCF₃, O-(C₁-C₆)alkyl, (C₁-C₄)alkoxyalkyl, S-(C₁-C₆)alkyl, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₃-C₈)cycloalkyl, O-(C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkenyl, O-(C₃-C₈)cycloalkenyl, (C₂-C₆)alkynyl, (C₀-C₈)alkylenearyl, -O-(C₀-C₈)alkylenearyl, S-aryl, N(R13)(R14), SO₂-CH₃, COOH, COO-(C₁-C₆)alkyl, CON(R15)(R16), N(R17)CO(R18), N(R19)SO₂(R20), CO(R21), or a 5- to 7-membered heterocycle having 1-4 heteroatoms;

15 R13, R14 are independently of one another H, (C₁-C₆)alkyl, or R13 and R14 together with the nitrogen atom to which they are bonded form a 5- to 6-membered ring, where, in the case of the 6-membered ring, a CH₂ group may be replaced by O or S;

20 R15, R16 are independently of one another H, (C₁-C₆)alkyl, or R15 and R16 together with the nitrogen atom to which they are bonded form a 5- to 6-membered ring, where, in the case of the 6-membered ring, a CH₂ group may be replaced by O or S;

25 R17, R19 are independently of one another H, or (C₁-C₆)alkyl;

R18, R20, R21 are independently of one another (C₁-C₆)alkyl, or aryl;

B is N(R24), or O;

30

R24 is H, or (C₁-C₆)alkyl;

R5 is H, or (C₁-C₆)alkyl;

W is N, or C(R25);

R25 is H, (C₁-C₆)alkyl, aryl, or a bond to Y;

5

T is N, or C(R26);

R26 is H, (C₁-C₆)alkyl, aryl, (C₀-C₈)alkylenearyl, or a bond to Y;

10 U is O, S, N(R27), -C(R30)=N-, or -N=C(R31)-;

wherein R27, R30, R31 are independently of one another H, (C₁-C₆)alkyl, a bond to Y;

15 Y is (C₁-C₈)alkylene, in which one or more carbons may be replaced by O, S, SO, SO₂, C(R32)(R33), CO, C(R34)(OR35) or N(R36);

R32, R33, R34, R35, R36 are independently of one another H, (C₁-C₆)alkyl, or aryl;

20 R6, R7 are independently of one another H, (C₁-C₆)alkyl, (C₃-C₇)cycloalkyl, or R6 and Y or R6 and R7 together with the nitrogen atom to which they are bonded form a 3- to 8-membered ring in which one or more carbons may be replaced by O, N or S and the 3- to 8-membered ring may carry further substituents, such as (C₁-C₆)alkyl, aryl, CON(R37)(R38), N(R39)(R40), OH, O-(C₁-
25 C₆)alkyl or NHCO(C₁-C₆)alkyl;

R37, R38, R39, R40 are independently of one another H, or (C₁-C₆)alkyl;

and the physiologically acceptable salts thereof.

30

Preference is given to compounds of formula I, in which one or more radicals have the following meaning:

A is (C₂-C₇)alkyl, (C₀-C₃)alkylenearyl; or a 4- to 10-membered mono- or bicyclic ring which may contain one or more heteroatoms selected from the group consisting of N, O and S, and the 4- to 10-membered ring may carry further substituents, such as F, Cl, Br, NO₂, CF₃, (C₁-C₆)alkyl, aryl, CON(R37)(R38),
5 N(R39)(R40), O-(C₁-C₆)alkyl, or NHCO(C₁-C₆)alkyl;

X is a bond, C(R8)(R9), O, N(R12), S, or SO₂;

10 R8, R9, R12 are independently of one another H, or (C₁-C₆)alkyl;

D is N, or C(R41);

E is N, or C(R42);

15 G is N, or C(R43);

L is N, or C(R44);

20 where the total number of the nitrogen atoms defined by D, E, G and L is 0, 1 or 2;

R1, R2, R3, R41, R42, R43, R44 are independently of one another H, F, Cl, Br, CF₃, NO₂, O-(C₁-C₆)alkyl, (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, O-(C₃-C₈)cycloalkyl, (C₂-C₆)alkynyl, (C₀-C₈)alkylenearyl, -O-(C₀-C₃)alkylenearyl, S-aryl, N(R13)(R14),
25 SO₂-CH₃, COO-(C₁-C₆)alkyl, CON(R15)(R16), N(R17)CO(R18), N(R19)SO₂(R20), or CO(R21);

30 R13, R14 are independently of one another H, (C₁-C₆)alkyl, or R13 and R14 together with the nitrogen atom to which they are bonded form a 5- to 6-membered ring, where, in the case of the 6-membered ring, a CH₂ group may be replaced by O or S;

R15, R16 are independently of one another H, (C₁-C₆)alkyl, or R15 and R16 together with the nitrogen atom to which they are bonded form a 5- to 6-membered ring, where, in the case of the 6-membered ring, a CH₂ group may be replaced by O or S;

5

R17, R19 are independently of one another H, or (C₁-C₆)alkyl;

R18, R20, R21 are independently of one another (C₁-C₆)alkyl, or aryl;

10

B is N(R24), or O;

R24 is H, or (C₁-C₆)alkyl;

R5 is H, or (C₁-C₆)alkyl;

15

W is N, or C(R25);

R25 is H, (C₁-C₆)alkyl, or aryl;

20

T is C(R26);

R26 is H, (C₁-C₆)alkyl, aryl, or a bond to Y;

U is O, S, N(R27), or -N=C(R31)-;

25

wherein R27, R31 are independently of one another H, (C₁-C₆)alkyl, or a bond to Y;

Y is (C₁-C₄)alkylene, in which a carbon may be replaced by SO₂, C(R32)(R33), CO or N(R36);

30

R32, R33, R36 are independently of one another H, (C₁-C₆)alkyl, or aryl;

R6, R7 are independently of one another H, (C₁-C₆)alkyl, (C₃-C₇)cycloalkyl, or R6 and Y or R6 and R7 together with the nitrogen atom to which they are bonded form a 4- to 7-membered ring in which one or more carbons may be replaced by O, N or S and the 4- to 7-membered ring may carry further
5 substituents such as (C₁-C₆)alkyl, aryl, CON(R37)(R38), N(R39)(R40), OH or NHCO(C₁-C₆)alkyl;

R37, R38, R39, R40 are independently of one another H, or (C₁-C₆)alkyl;
10 and the physiologically acceptable salts thereof.

Particular preference is given to compounds of formula I, in which one or more radicals have the following meaning:

15 A is (C₃-C₇)alkyl, (C₀-C₂)alkylenearyl; a 5- to 10-membered mono- or bicyclic ring which may contain 0, 1 or 2 heteroatoms selected from the group consisting of N, O and S, and the 5- to 10-membered ring may carry further substituents, such as F, Cl, Br, NO₂, CF₃, (C₁-C₆)alkyl, aryl, O-(C₁-C₆)alkyl or NHCO(C₁-C₆)alkyl;

20 X is a bond, C(R8)(R9), O, or N(R12);

R8, R9, R12 are independently of one another H, or (C₁-C₆)alkyl;

25 D is N, or C(R41);

E is N, or C(R42);

G is N, or C(R43);
30

L is N, or C(R44);

where the total number of the nitrogen atoms defined by D, E, G and L is 0 or 1;

5 R1, R2, R3, R41, R42, R43, R44 are independently of one another H, F, Cl, CF₃, NO₂, O-(C₁-C₆)alkyl, (C₁-C₆)alkyl, O-(C₃-C₈)cycloalkyl, (C₀-C₂)alkylenearyl, -O-(C₀-C₃)alkylenearyl, N(R13)(R14), COO-(C₁-C₆)alkyl, CON(R15)(R16), N(R17)CO(R18), N(R19)SO₂(R20), or CO(R21);

10 R13, R14 are independently of one another H, or (C₁-C₆)alkyl,

R15, R16 are independently of one another H, or (C₁-C₆)alkyl,

R17, R19 are independently of one another H, or (C₁-C₆)alkyl;

15 R18, R20, R21 are independently of one another (C₁-C₆)alkyl, or aryl;

B is N(R24);

20 R24 is H, or (C₁-C₆)alkyl;

R5 is H, or (C₁-C₆)alkyl;

W is N, or C(R25);

25 R25 is H, or (C₁-C₆)alkyl;

T is C(R26);

30 R26 is H, (C₁-C₆)alkyl, or a bond to Y;

U is O, S, or N(R27);

R27 is H, (C₁-C₆)alkyl, or a bond to Y;

Y is (C₁-C₃)alkylene, in which a carbon may be replaced by SO₂, C(R₃₂)(R₃₃) or CO;

5 R₃₂, R₃₃ are independently of one another H, (C₁-C₆)alkyl, or aryl;

R₆, R₇ are independently of one another H, (C₁-C₆)alkyl, (C₃-C₇)cycloalkyl, or R₆ and Y or R₆ and R₇ together with the nitrogen to which they are bonded form a 5- to or 6-membered ring in which one or more carbons may be
10 replaced by O or N and the 5- or 6-membered ring may carry further substituents, such as (C₁-C₆)alkyl, aryl, CON(R₃₇)(R₃₈), N(R₃₉)(R₄₀), OH or NHCO(C₁-C₆)alkyl;

R₃₇, R₃₈, R₃₉, R₄₀ are independently of one another H, or (C₁-C₆)alkyl;
15 and the physiologically acceptable salts thereof.

The invention relates to compounds of formula I in the form of their racemates, enantiomer-enriched mixtures and pure enantiomers and to their
20 diastereomers and mixtures thereof.

The substituents R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₅, R₂₆, R₂₇, R₃₀, R₃₁, R₃₂, R₃₃, R₃₄, R₃₅, R₃₆, R₃₇, R₃₈, R₃₉, R₄₀, R₄₁, R₄₂, R₄₃ and R₄₄ may have
25 straight-chain, branched or optionally halogenated alkyl, alkylene, alkenyl and alkynyl radicals.

The term "aryl" means a phenyl or naphthyl group. The term "ring" means a cyclic structure which may be aromatic, partly saturated or completely saturated.
30 The optional ring formation of R₆, Y and the nitrogen to which they are bonded can be illustrated by examples 6 and 16 without limiting the general description mentioned above.

Pharmaceutically acceptable salts are particularly suitable for medical applications, due to their greater solubility in water compared with the starting or base compounds. Said salts must have a pharmaceutically acceptable anion or cation. Suitable pharmaceutically acceptable acid addition salts of the compounds of the invention are salts of inorganic acids, such as hydrochloric acid, hydrobromic acid, phosphoric acid, metaphosphoric acid, nitric acid, sulfonic acid and sulfuric acid and also of organic acids, such as, for example, acetic acid, benzenesulfonic acid, benzoic acid, citric acid, ethanesulfonic acid, fumaric acid, gluconic acid, glycolic acid, isethionic acid, lactic acid, lactobionic acid, maleic acid, malic acid, methanesulfonic acid, succinic acid, p-toluenesulfonic acid, tartaric acid and trifluoroacetic acid. For medicinal purposes, particular preference is given to using the chloride salt. Suitable pharmaceutically acceptable basic salts are ammonium salts, alkali metal salts (such as sodium salts and potassium salts) and alkaline earth metal salts (such as magnesium salts and calcium salts).

Salts having a pharmaceutically unacceptable anion are likewise included within the scope of the present invention as useful intermediates for preparing or purifying pharmaceutically acceptable salts and/or for use in nontherapeutic applications, for example *in-vitro* applications.

The term "physiologically functional derivative" used herein relates to any physiologically acceptable derivative of an inventive compound of formula I, for example, an ester which on administration to a mammal (e.g., humans) is capable of forming (directly or indirectly) a compound of formula I or an active metabolite thereof.

The physiologically functional derivatives also include prodrugs of the compounds of the invention. Such prodrugs may be metabolized *in vivo* to a compound of the invention. These prodrugs may or may not be active themselves.

The compounds of the invention may also be present in various polymorphous forms, for example as amorphous and crystalline polymorphous

forms. All polymorphous forms of the compounds of the invention are included within the scope of the invention and are another aspect of the invention.

5 All references to "compound(s) according to formula (I)" refer hereinbelow to a compound/compounds of the formula (I) as described above and also to their salts, solvates and physiologically functional derivatives as described herein.

10 The amount of a compound according to formula (I) which is required in order to attain the desired biological effect depends on a number of factors, for example the specific compound selected, the intended use, the type of administration and the clinical state of the patient. In general, the daily dose is in the range from 0.3 mg to 100 mg (typically from 3 mg to 50 mg) per day per kilogram of body weight, for example 3-10 mg/kg/day. An intravenous dose can be, for example, in the range from 0.3 mg to 1.0 mg/kg and can be administered in
15 a suitable manner as an infusion of 10 ng to 100 ng per kilogram per minute. Suitable infusion solutions for these purposes may contain, for example, from 0.1 ng to 10 mg, typically from 1 ng to 10 mg per milliliter. Individual doses may contain, for example, from 1 mg to 10 g of the active compound. Thus, ampoules for injections can contain, for example, from 1 mg to 100 mg, and orally
20 administerable individual dose formulations such as, for example, tablets or capsules can contain, for example, from 1.0 to 1000 mg, typically from 10 to 600 mg. In the case of pharmaceutically acceptable salts, the abovementioned masses relate to the mass of the free compound on which the salt is based. The compound used for the prophylaxis or therapy of the abovementioned conditions
25 may be the compounds according to formula (I) themselves, but they are preferably present in the form of a pharmaceutical composition together with an acceptable carrier. The carrier must be naturally acceptable, in the sense that it is compatible with the other ingredients of said composition and is not harmful to the patient's health. The carrier may be a solid or a liquid or both and is preferably
30 formulated with the compound as an individual dose, for example, as a tablet which may contain from 0.05% to 95% by weight of the active compound. Further pharmaceutically active substances may also be present, including further compounds according to formula (I). The pharmaceutical compositions of the

invention may be prepared according to any of the known pharmaceutical methods which essentially comprise mixing the ingredients with pharmacologically acceptable carriers and/or excipients.

5 Pharmaceutical compositions of the invention are those which are suitable for oral, rectal, topical, peroral (e.g., sublingual) and parenteral (e.g., subcutaneous, intramuscular, intradermal or intravenous) administration, although the most suitable manner of administration depends in each individual case on the nature and severity of the condition to be treated and on the nature of the
10 compound according to formula (I) used in each case. Sugar-coated formulations and sugar-coated delayed-release formulations, too, are included within the scope of the invention. Preference is given to acid-resistant and enteric formulations. Suitable enteric coatings include cellulose acetate phthalate, polyvinyl acetate phthalate, hydroxypropylmethylcellulose phthalate and anionic polymers of
15 methacrylic acid and methyl methacrylate.

 Suitable pharmaceutical compounds for oral administration may be present in separate units as, for example, capsules, cachets, lozenges or tablets, which in each case contain a particular amount of the compound according to formula (I);
20 as powders or granules; as solution or suspension in an aqueous or nonaqueous liquid; or as an oil-in-water or water-in-oil emulsion. As already mentioned, said compositions can be prepared according to any suitable pharmaceutical method which includes a step in which the active compound and the carrier (which may comprise one or more additional components) are contacted. In general, the
25 compositions are prepared by uniform and homogeneous mixing of the active compound with a liquid and/or finely dispersed solid carrier, after which the product is shaped, if necessary. Thus, a tablet, for example, may be prepared by pressing or shaping a powder or granules of the compound, where appropriate with one or more additional components. Pressed tablets can be prepared by
30 tableting the compound in free-flowing form, for example, a powder or granules, mixed, where appropriate, with a binder, lubricant, inert diluent and/or one or more surface active/dispersing agents in a suitable machine. Shaped tablets can be

prepared by shaping the pulverulent compound, moistened with an inert liquid diluent, in a suitable machine.

5 Pharmaceutical compositions which are suitable for peroral (sublingual) administration include lozenges which contain a compound according to formula (I) with a flavoring, usually sucrose and gum arabic or tragacanth, and pastilles which comprise the compound in an inert base such as gelatin and glycerol or sucrose and gum arabic.

10 Suitable pharmaceutical compositions for parenteral administration preferably comprise sterile aqueous preparations of a compound according to formula (I) which are preferably isotonic with the blood of the intended recipient. These preparations are preferably administered intravenously, although they may also be administered subcutaneously, intramuscularly or intradermally as an
15 injection. Said preparations may preferably be prepared by mixing the compound with water and rendering the obtained solution sterile and isotonic with the blood. Injectable compositions of the invention generally contain from 0.1 to 5% by weight of the active compound.

20 Suitable pharmaceutical compositions for rectal administration are preferably present as individual dose suppositories. These may be prepared by mixing a compound according to formula (I) with one or more conventional solid carriers, for example, cocoa butter, and shaping the resulting mixture.

25 Suitable pharmaceutical compositions for topical application to the skin are preferably present as ointment, cream, lotion, paste, spray, aerosol or oil. Carriers which may be used are petroleum jelly, lanolin, polyethylene glycols, alcohols and combinations of two or more of these substances. In general, the active compound is present at a concentration of from 0.1 to 15%, for example from 0.5 to 2%, by
30 weight of the composition.

Transdermal administration is also possible. Suitable pharmaceutical compositions for transdermal administration may be present as individual patches

which are suitable for long-term close contact with the epidermis of the patient. Such patches suitably contain the active compound in an optionally buffered aqueous solution, dissolved and/or dispersed in an adhesive or dispersed in a polymer. A suitable active compound concentration is from approx. 1% to 35%,
5 preferably approx. 3% to 15%. A particular possibility is the release of the active compound by electrotransport or iontophoresis, as described, for example, in *Pharmaceutical Research*, 2(6):318 (1986).

The compounds of formula I are distinguished by beneficial actions on the
10 metabolism of lipids, and they are particularly suitable for weight reduction and, after weight reduction, for maintaining a reduced weight in mammals and as anorectic agents. The compounds are distinguished by their low toxicity and their few side effects. The compounds may be employed alone or in combination with other weight-reducing or anorectic active compounds. Further anorectic active
15 compounds of this kind are mentioned, for example, in the Rote Liste 2001, Arzneimittelverzeichnis für Deutschland, Rote Liste Service GmbH, Frankfurt, under weight-reducing agents/appetite suppressants, and may also include those active compounds which increase the energy turnover of the organism and thus lead to weight reduction or else those which influence the general metabolism of
20 said organism such that increased calorie intake does not cause an enlargement of the fat depots and a normal calorie intake causes a reduction in the fat depots of said organism. The compounds are suitable for the prophylaxis and, in particular, for the treatment of problems of excess weight or obesity. The compounds are furthermore suitable for the prophylaxis and, in particular, for the
25 treatment of type II diabetes, of arteriosclerosis and for the normalization of lipid metabolism and for the treatment of high blood pressure. The compounds act as MCH antagonists and are also suitable for the treatment of paresthesia and other psychiatric indications such as, for example, depressions, anxieties, anxiety neuroses, schizophrenia and also for the treatment of disorders associated with
30 the circadian rhythm and for the treatment of drug abuse.

In a further aspect of the invention, the compounds of formula I may be administered in combination with one or more further pharmacologically active substances which may be selected, for example, from the group consisting of

antidiabetics, antiadipose agents, blood-pressure-lowering active compounds, lipid reducers and active compounds for the treatment and/or prevention of complications caused by diabetes or associated with diabetes.

Suitable antidiabetics include insulins, amylin, GLP-1 and GLP-2 derivatives
5 such as, for example, those disclosed by Novo Nordisk A/S in WO 98/08871 and also oral hypoglycemic active compounds.

Said oral hypoglycemic active compounds preferably include sulfonyl ureas, biguanidines, meglitinides, oxadiazolidinediones, thiazolidinediones, glucosidase inhibitors, glucagon receptor antagonists, GLP-1 agonists, potassium channel
10 openers such as, for example, those disclosed by Novo Nordisk A/S in WO 97/26265 and WO 99/03861, insulin sensitizers, activators of insulin receptor kinase, inhibitors of liver enzymes involved in the stimulation of gluconeogenesis and/or glycogenolysis, for example glycogen phosphorase inhibitors, modulators of glucose uptake and glucose elimination, lipid metabolism-modifying compounds
15 such as antihyperlipidemic active compounds and antilipidemic active compounds, for example HMGCoA-reductase inhibitors, inhibitors of cholesterol transport/cholesterol uptake, inhibitors of the reabsorption of bile acid or inhibitors of microsomal triglyceride transfer protein (MTP), compounds which reduce food intake, PPAR and RXR agonists and active compounds which act on the ATP-
20 dependent potassium channel of beta cells.

In one embodiment of the present invention, the present compounds are administered in combination with insulin.

In another embodiment, the compounds of the invention are administered in combination with a sulfonylurea such as, for example, tolbutamide, glibenclamide,
25 glimepiride, glipizide, gliquidone, glisoxepide, glibornuride or gliclazide.

In another embodiment, the compounds of the present invention are administered in combination with a biguanidine such as, for example, metformin.

In another embodiment, the compounds of the present invention are administered in combination with a meglitinide such as, for example, repaglinide.

In yet another embodiment, the compounds of the present invention are administered in combination with a thiazolidinedione such as, for example,
5 troglitazone, ciglitazone, pioglitazone, rosiglitazone or the compounds disclosed by Dr. Reddy's Research Foundation in WO 97/41097, in particular 5-[[4-[(3,4-dihydro-3-methyl-4-oxo-2-quinazolinylmethoxy)phenyl]methyl]-2,4-thiazolidinedione.

In another embodiment, the compounds of the present invention are administered in combination with an α -glucosidase inhibitor such as, for example,
10 miglitol or acarbose.

In another embodiment, the compounds of the present invention are administered in combination with an active compound which acts on the ATP-dependent potassium channel of the beta cells, such as, for example, tolbutamide, glibenclamide, glimepiride, glipizide, gliclazide or repaglinide.

15 In yet another embodiment, the compounds of the present invention are administered in combination with an antihyperlipidemic active compound or an antilipidemic active compound such as, for example, cholestyramine, colestipol, clofibrate, fenofibrate, gemfibrozil, lovastatin, pravastatin, simvastatin, atorvastatin, cerivastatin, fluvastatin, probucol, ezetimibe or dextrothyroxine.

20 In another embodiment, the compounds of the present invention are administered in combination with more than one of the aforementioned compounds, for example in combination with a sulfonylurea and metformin, a sulfonylurea and acarbose, repaglinide and metformin, insulin and a sulfonylurea, insulin and metformin, insulin and troglitazone, insulin and lovastatin, etc.

25 Furthermore, the compounds of the invention may be administered in combination with one or more antiadipose agents or appetite-controlling active compounds.

Such active compounds may be selected from the group consisting of
CART agonists, NPY antagonists, MC4 agonists, orexin antagonists, H3 agonists,
TNF agonists, CRF agonists, CRF BP antagonists, urocortin agonists, β 3 agonists,
MSH (melanocyte-stimulating hormone) agonists, CCK agonists, serotonin re-
5 uptake inhibitors, mixed serotonin and noradrenalin reuptake inhibitors, 5HT
modulators, MAO inhibitors, bombesin agonists, galanin antagonists, growth
hormone, growth-hormone-releasing compounds, TRH agonists, uncoupling
protein 2 or 3 modulators, leptin agonists, dopamine agonists (bromocriptine,
doprexin), lipase/amylase inhibitors, cannabinoid receptor 1 antagonists,
10 modulators of acylation-stimulating protein (ASP), PPAR modulators, RXR
modulators, hCNTF mimetics or TR- β agonists.

In one embodiment of the invention, the antiadipose agent is leptin or
modified leptin.

In another embodiment, the antiadipose agent is dexamphetamine or
15 amphetamine.

In another embodiment, the antiadipose agent is fenfluramine or
dexfenfluramine.

In yet another embodiment, the antiadipose agent is sibutramine or the
mono- and bis-demethylated active metabolite of sibutramine.

20 In another embodiment, the antiadipose agent is orlistate.

In another embodiment, the antiadipose agent is mazindol, diethylpropione
or phentermine.

Furthermore, the compounds of the present invention may be administered
in combination with one or more antihypertensive active compounds. Examples of
25 antihypertensive active compounds are beta blockers such as alprenolol, atenol,
timolol, pindolol, propanolol and metoprolol, ACE (angiotensin-converting enzyme)
inhibitors such as, for example, benazepril, captopril, enalapril, fosinopril, lisinopril,

quinapril and rampril, calcium channel blockers such as nifedipine, felodipine, nicardipine, isradipine, nimodipine, diltiazem and verapamil, and also alpha blockers such as doxazosin, urapidil, prazosin and terazosin. Furthermore, reference may be made to Remington: The Science and Practice of Pharmacy,
5 19th edition, Gennaro, editor, Mack Publishing Co., Easton, PA, 1995.

It is self-evident that every suitable combination of the compounds of the invention with one or more of the aforementioned compounds and optionally one or more other pharmacologically active substances is to be regarded as covered by the scope of protection of the present invention.

EXAMPLES

The activity of the compounds was assayed as follows:

Biological test model:

The anorectic action was tested on female NMRI mice. After removal of feed for 17 hours, the preparation to be tested was administered by gavage. The animals were housed singly and, with free access to drinking water, they were offered evaporated milk 30 minutes after administration of the preparation. The consumption of evaporated milk was determined and the general behavior of the animals were monitored every half an hour for 7 hours. The measured milk consumption was compared to that of vehicle-treated control animals.

Table 1: Anorectic action, measured as a reduction in the cumulative milk consumption by treated animals compared with control animals

Example	Oral dose [mg/kg]	Number of animals/ cumulative milk consumption by treated animals N / [mL]	Number of animals/ cumulative milk consumption by control animals N / [mL]	Reduction in cumulative milk consumption as % of the control
Example 1	30	5/2.28	5/3.26	30
Example 4	10	5/2.74	5/4.44	38

The table indicates that the compounds of formula I exhibit very good anorectic action.

In two simultaneously published articles in *Nature* (*Nature*, 400:261-264, 1999; *Nature*, 400:265-269, 1999, see enclosure), two groups separately described a highly specific receptor for melanin-concentrating hormone (MCH). MCH takes over important functions in the control of food intake. Compounds

acting on the MCH receptor therefore have anorectic action and are suitable for the treatment of obesity. The test for anorectic action of the inventive compounds of formula I was therefore carried out as follows.

5 Functional measurements for determination of IC50

Cloning of the cDNA for human MCH receptor, preparation of a recombinant HEK293 cell line expressing human MCH receptor and functional measurements with said recombinant cell line were carried out according to the description by Audinot *et al.* (*J. Biol. Chem.*, 276, 13554-13562, 2001). In contrast
10 to the reference, however, plasmid pEAK8 from EDGE Biosystems (USA) was used for constructing the expression vector. A transformed HEK cell line named "PEAK Stable Cells" (likewise from EDGE Biosystems) served as host for transfection. The functional measurements of cellular calcium flow, after addition
15 of agonists (MCH), in the presence of the ligand of the invention was carried out with the aid of the FLIPR instrument from Molecular Devices (USA), using the manufacturer's protocols.


Table 2: Test for anorectic action of the inventive compounds of formula I; results from the cellular assay

Example	IC50 / μ M	Example	IC50 / μ M	Example	IC50 / μ M
1	0.15	16	0.33	76	4.25
2	0.15	17	2.14	77	0.70
3	0.29	18	1.04	78	2.75
4	0.13	19	0.70	79	2.13
5	0.50	22	4.42	80	3.36
6	2.34	24	0.86	81	2.69
7	0.45	26	0.92	84	0.40
8	1.90	29	2.91	86	2.78
9	0.10	33	1.24	105	1.0

10	0.11	63	0.57	106	0.20
11	0.14	65	0.50	107	1.0
13	2.50	71	2.65	108	0.43
14	0.30	72	0.32	109	1.29
15	0.18	73	0.14		


The examples and preparation methods listed below serve to illustrate the invention but without limiting it.

Example 1 1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea




Carbonyldiimidazole (5.12 g) was added to a solution cooled to 0°C of 1-dimethylaminoethyl-5-aminoindole (6.30 g) in dimethylformamide (50 mL). After 10 minutes, 4-aminodiphenyl ether (5.84 g) was added and the reaction mixture was heated to 80°C for 2 hours. After cooling, the reaction was diluted with ethyl acetate and washed with water. The organic phase was dried over magnesium sulfate, filtered and concentrated. The residue was purified by chromatography on silica gel (eluent: dichloromethane/methanol 9:1). Thus the product having a molecular weight of 414.15 (C₂₅H₂₆N₄O₂); MS (ESI): 415 (M+H⁺) was obtained.

Example 2 1-(4-Butoxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea



The compound was prepared from 4-butoxyaniline and 1-
5 dimethylaminoethyl-5-aminoindole, as described in Example 1. Thus, the product
having a molecular weight of 394.52 ($C_{23}H_{30}N_4O_2$); MS (ESI): 395 ($M+H^+$) was
obtained.

Example 3 1-(1-Methyl-2-pyrrolidin-1-ylmethyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea



The compound was prepared from 4-aminodiphenyl ether and 1-methyl-2-
pyrrolidin-1-ylmethyl-1H-indol-5-ylamine, as described in Example 1. Thus, the
product having a molecular weight of 440.55 ($C_{27}H_{28}N_4O_2$); MS (ESI): 441 ($M+H^+$)
was obtained.

1-Methyl-2-pyrrolidin-1-ylmethyl-1H-indol-5-ylamine

Formic acid (0.11 mL) was added to a suspension of 1-methyl-5-nitro-2-
pyrrolidin-1-ylmethyl-1H-indole (150 mg), ethanol (2 mL) and palladium(II)
hydroxide on carbon (20%, 30 mg) and the suspension was heated to 60°C for 5
minutes. After gas production had ceased, the suspension was stirred for another
20 minutes and the catalyst was filtered off. The filtrate was concentrated and
distributed between saturated sodium carbonate solution and methyl tert-butyl
ether. The organic phase was removed, dried over magnesium sulfate and

concentrated. Thus, the product having a molecular weight of 229.33 ($C_{14}H_{19}N_3$); MS (ESI): 230 ($M+H^+$) was obtained.

1-Methyl-5-nitro-2-pyrrolidin-1-ylmethyl-1H-indole

Mesyl chloride (92 mg) was added dropwise to a solution cooled to 0°C of
5 (1-methyl-5-nitro-1H-indol-2-yl)methanol (121 mg) in dichloromethane (10 mL) and triethylamine (0.17 mL). After 15 minutes, pyrrolidine (142 mg) was added and the solution was then stirred at room temperature for 1 hour. The reaction solution was washed with saturated sodium carbonate solution, dried over magnesium sulfate and concentrated. The residue was purified via chromatography on silica
10 gel (eluent: ethyl acetate/triethylamine 99:1). Thus, the product having a molecular weight of 259.31 ($C_{14}H_{17}N_3O_2$); MS (ESI): 260 ($M+H^+$) was obtained.


(1-Methyl-5-nitro-1H-indol-2-yl)methanol

Sulfuric acid (96% strength, 0.64 mL) was added dropwise to a suspension cooled to 0°C of lithium aluminum hydride in tetrahydrofuran (50 mL) within 20
15 minutes. After 20 minutes, a solution of ethyl 1-methyl-5-nitro-1H-indole 2-carboxylate (1.85 g) in tetrahydrofuran (40 mL) was added dropwise. After 30 minutes, water (2 mL) was added. After 30 minutes, the resulting precipitate was filtered off and the filtrate was concentrated. The crude product was purified via chromatography on silica gel (eluent: n-heptane/ethyl acetate 3:2). Thus, the
20 product having a molecular weight of 206.20 ($C_{10}H_{10}N_2O_3$); MS (ESI): 207 ($M+H^+$) was obtained.

Ethyl 1-methyl-5-nitro-1H-indole 2-carboxylate

A suspension of ethyl 5-nitro-1H-indole 2-carboxylate (2.34 g), potassium carbonate (3.45 g), methyl iodide (2.13 g) and acetonitrile (30 mL) was kept at
25 60°C for 6 hours. After cooling to room temperature, water was added and the precipitated product was isolated by filtration. Thus, the product having a molecular weight of 248.24 ($C_{12}H_{12}N_2O_4$); MS (ESI): 249 ($M+H^+$) was obtained.

Example 4 1-[1-(2-Dimethylaminoethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea



Zinc dust (250 mg) was added to a solution of 1-[4-(2-
5 dimethylaminoethylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea (50 mg) in
dichloromethane (10 mL) and glacial acetic acid (1 mL). After 10 minutes, the
inorganic material was filtered off via kieselguhr. The filtrate was washed with a
sodium carbonate solution (10% strength), dried over magnesium sulfate and
concentrated. The residue was taken up in dichloromethane (5 mL) and ethanol
10 (5 mL) and admixed with dimethylformamide dimethyl acetal (0.3 mL) and formic
acid (0.3 mL). Dichloromethane was evaporated by heating the mixture by means
of a hot-air gun. The remaining mixture was concentrated and distributed between
dichloromethane and a sodium carbonate solution (10% strength). The organic
phase was removed, dried and concentrated. The residue was purified by
15 preparative HPLC. Thus, the product having a molecular weight of 415.50
(C₂₄H₂₅N₅O₂); MS (ESI): 416 (M+H⁺) was obtained. Melting point of the
hydrochloride: 213-215 °C.

1-[4-(2-Dimethylaminoethylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea

A solution of 2-dimethylaminoethylamine in dimethylformamide (1M, 2 mL)
20 and 1-(4-fluoro-3-nitrophenyl)-3-(4-phenoxyphenyl)urea (200 mg) was stirred for
48 hours. The mixture was distributed between dichloromethane and a sodium
carbonate solution (10% strength). The organic phase was dried and
concentrated. The residue was recrystallized from toluene. Melting point:
178-180°C.


25 1-(4-Fluoro-3-nitrophenyl)-3-(4-phenoxyphenyl)urea

4-Fluoro-3-nitrophenyl isocyanate (2.2 mmol) was added to a solution of
4-phenoxyaniline (2 mmol) in dimethylformamide (20 mL). After 2 days, the

reaction mixture was distributed between dichloromethane and a saturated sodium carbonate solution. The organic phase was dried and concentrated. The residue was purified via chromatography on silica gel (eluent: ethyl acetate/dichloromethane 95:5) and subsequent recrystallization from ethyl acetate/hexane.

5 Melting point: 174-176°C.

Example 5 1-[1-(2-Dimethylaminoethyl)-2-methyl-1H-benzimidazol-5-yl]-3-(4-isopropoxyphenyl)urea



10 1-[4-(2-Dimethylaminoethylamino)-3-nitrophenyl]-3-(4-isopropoxyphenyl)urea (75 mg) was reduced using zinc dust, as described in Example 4. The reaction product was dissolved in methanol and admixed with triethyl orthoacetate (0.5 mL) and glacial acetic acid (0.2 mL). The mixture was heated under reflux for 5 minutes. Volatile components were removed. The
15 residue was distributed between dichloromethane and a sodium carbonate solution. The organic phase was dried and concentrated. The residue was purified by preparative HPLC. Thus, the product having a molecular weight of 395.51 (C₂₂H₂₉N₅O₂); MS (ESI): 396 (M+H⁺) was obtained.


1-[4-(2-Dimethylaminoethylamino)-3-nitrophenyl]-3-(4-isopropoxyphenyl)urea

20 The compound was obtained from 1-(4-fluoro-3-nitrophenyl)-3-(4-isopropoxyphenyl)urea and 2-dimethylaminoethylamine as in Example 4. The compound was reacted further without purification.

1-(4-Fluoro-3-nitrophenyl)-3-(4-isopropoxyphenyl)urea

The compound was obtained from 4-fluoro-3-nitrophenyl isocyanate and 4-isopropoxyaniline as in Example 4. Melting point: 170-172°C.

Example 6 1-[1-(1-Ethylpyrrolidin-2-ylmethyl)-2-methyl-1H-benzoimidazol-5-yl]-3-(4-isopropoxyphenyl)urea




The compound was prepared from 1-{4-[(1-ethylpyrrolidin-2-ylmethyl)amino]-3-nitrophenyl}-3-(4-isopropoxyphenyl)urea, as described in Example 5. Thus, the product having a molecular weight of 435.57 ($C_{25}H_{33}N_5O_2$); MS (ESI): 436 ($M+H^+$) was obtained. Melting point: (ethyl acetate/hexane): 185-187°C.

1-{4-[(1-Ethylpyrrolidin-2-ylmethyl)amino]-3-nitrophenyl}-3-(4-isopropoxyphenyl)urea

The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-isopropoxyphenyl)urea and 1-ethylpyrrolidin-2-ylmethylamine, as described in Example 4, and reacted further without any further purification.

Example 7 1-(4-Isopropoxyphenyl)-3-[2-methyl-1-(2-piperidin-1-ylethyl)-1H-benzoimidazol-5-yl]urea




The compound was prepared from 1-(4-isopropoxyphenyl)-3-[3-nitro-4-(2-piperidin-1-yl-ethylamino)phenyl]urea, as described in Example 5. Thus the product having a molecular weight of 435.57 ($C_{25}H_{33}N_5O_2$); MS (ESI): 436 ($M+H^+$) was obtained.

5 1-(4-Isopropoxyphenyl)-3-[3-nitro-4-(2-piperidin-1-ylethylamino)phenyl]urea

The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-isopropoxyphenyl)urea and 1-(2-aminoethyl)piperidine (60°C, 4 h), as described in Example 4. Melting point (ethyl acetate): 157-159°C.

10 **Example 8 1-(4-Isopropoxyphenyl)-3-[2-methyl-1-(2-morpholin-4-yl-ethyl)-1H-benzoimidazol-5-yl]urea**




15 The compound was prepared from 1-(4-isopropoxyphenyl)-3-[4-(2-morpholin-4-ylethylamino)-3-nitrophenyl]urea, as described in Example 5. Thus, the product having a molecular weight of 437.55 ($C_{24}H_{31}N_5O_3$); MS (ESI): 438 ($M+H^+$) was obtained.

1-(4-Isopropoxyphenyl)-3-[4-(2-morpholin-4-yl-ethylamino)-3-nitrophenyl]urea

The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-isopropoxyphenyl)urea and 1-(2-aminoethyl)morpholine (60°C, 4 h), as described in Example 4. Melting point (ethyl acetate): 191-193°C.

20

Example 9 1-(4-Isopropoxyphenyl)-3-[2-methyl-1-(2-pyrrolidin-1-ylethyl)-1H-benzoimidazol-5-yl]urea




The compound was prepared from 1-[3-nitro-4-(2-pyrrolidin-1-ylethylamino)-phenyl]-3-(4-phenoxyphenyl)urea, as described in Example 5. Thus, the product having a molecular weight of 455.56 ($C_{27}H_{29}N_5O_2$); MS (ESI): 456 ($M+H^+$) was obtained.

1-[3-Nitro-4-(2-pyrrolidin-1-ylethylamino)phenyl]-3-(4-phenoxyphenyl)urea


The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-phenoxyphenyl)urea and 1-(2-aminoethyl)pyrrolidine (60°C, 5 h), as described in Example 4. Melting point (ethyl acetate/hexane): 179-181°C.

Example 10 1-[2-Methyl-1-(2-dimethylaminoethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea




The compound was prepared from 1-[4-(2-dimethylaminoethylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea, as described in Example 5. Thus, the product having a molecular weight of 429.53 ($C_{25}H_{27}N_5O_2$); MS (ESI): 430 ($M+H^+$) was obtained.

Example 11 1-(4-Phenoxyphenyl)-3-[1-(2-pyrrolidin-1-ylethyl)-1H-benzoimidazol-5-yl]urea




The compound was prepared from 1-[3-nitro-4-(2-pyrrolidin-1-ylethylamino)-phenyl]-3-(4-phenoxyphenyl)urea, as described in Example 4. Thus, the product having a molecular weight of 441.54 ($C_{26}H_{27}N_5O_2$); MS (ESI): 442 ($M+H^+$) was obtained.

Example 12 1-[2-Benzyl-1-(2-dimethylaminoethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea




1-[4-(2-Dimethylaminoethylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea (75 mg) was reduced as described in Example 4. The crude product was treated with phenylacetic acid (0.33 mmol), activated with HATU (0.33 mmol), and diisopropylamine (0.7 mmol) in dimethylformamide (1.5 mL) for 3 hours. The reaction mixture was distributed between dichloromethane and a sodium carbonate solution (10% strength). The organic phase was dried and concentrated. The residue was heated under reflux in trifluoroacetic acid (1 mL), water (1 mL) and acetonitrile (0.5 mL) for 5 minutes. Volatile components were evaporated and the residue was purified by preparative HPLC. Thus, the product having a molecular weight of 505.63 ($C_{31}H_{31}N_5O_2$); MS (ESI): 506 ($M+H^+$) was obtained.

Example 13 1-[1-(2-Dimethylaminoethyl)-2-phenyl-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea



1-[4-(2-Dimethylaminoethylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea
5 (50 mg) was reduced as described in Example 4. After filtration via kieselguhr, benzaldehyde (0.2 mL) was added to the filtrate. The reaction mixture was washed with a sodium carbonate solution (10% strength), dried and admixed with manganese dioxide (0.5 g). After 15 minutes, the inorganic material was filtered off and the filtrate was concentrated. The crude product was purified by
10 preparative HPLC. Thus, the product having a molecular weight of 491.60 ($C_{30}H_{29}N_5O_2$); MS (ESI): 492 ($M+H^+$) was obtained.


Example 14 1-[2-Ethyl-1-(2-pyrrolidin-1-ylethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea



15 1-[4-(2-Pyrrolidinoethylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea was reduced as described in Example 4. The crude product was reacted with triethyl orthopropionate according to Example 5. The crude product was purified by preparative HPLC. Thus, the product having a molecular weight of 469.59 ($C_{28}H_{31}N_5O_2$); MS (ESI): 470 ($M+H^+$) was obtained.

20

Example 15 1-[2-Methyl-1-(2-piperidin-1-ylethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea




1-[2-Methyl-1-(2-piperidin-1-ylethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea was reduced as described in Example 4. The crude product was reacted with triethyl orthoacetate, as described in Example 5. The crude product was purified by preparative HPLC. Thus, the product having a molecular weight of 469.59 ($C_{28}H_{31}N_5O_2$); MS (ESI): 470 ($M+H^+$) was obtained.

1-[2-Methyl-1-(2-piperidin-1-ylethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)-urea

The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-phenoxyphenyl)urea and 1-(2-aminoethyl)piperidine (60°C, 4 h), as described in Example 4. Melting point (ethyl acetate/hexane): 163-165°C.

Example 16 1-[1-(1-Ethylpyrrolidin-2-ylmethyl)-2-methyl-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea




1-[4-[(1-Ethylpyrrolidin-2-ylmethyl)amino]-3-nitrophenyl]-3-(4-phenoxyphenyl)urea was reduced as described in Example 4. The crude product was reacted with triethylorthoacetate, as described in Example 5. The crude product was purified by preparative HPLC. Thus, the product having a molecular weight of 469.59 ($C_{28}H_{31}N_5O_2$); MS (ESI): 470 ($M+H^+$) was obtained.

1-[4-[(1-Ethylpyrrolidin-2-yl)methyl]amino]-3-nitrophenyl]-3-(4-phenoxyphenyl)urea

The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-phenoxyphenyl)urea and C-(1-ethylpyrrolidin-2-yl)methylamine (60°C, 4 h), as described in Example 4. Melting point (ethyl acetate/hexane): 129-132°C.

5 **Example 17 1-(2-Dimethylaminomethyl-1H-benzoimidazol-5-yl)-3-(4-phenoxyphenyl)urea**




1-[4-(2,4-Dimethoxybenzylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea (75 mg) was reduced as described in Example 4. The reduced product was
10 reacted with dimethylaminoacetic acid (1 mmol), HATU (1 mmol) and diisopropylamine (2 mmol) in dimethylformamide (3 mL). After 3 hours, the mixture was distributed between ethyl acetate and a sodium carbonate solution. The organic phase was dried and concentrated. The crude product was purified by preparative HPLC. Thus the intermediate (N-{2-amino-5-[3-(4-
15 phenoxyphenyl)ureido]phenyl}-2-dimethylaminoacetamide) having a molecular weight of 419.49 (C₂₃H₂₅N₅O₃); MS (ESI): 420 (M+H⁺) was obtained.

This material was heated under reflux with pivalic acid and volatile components were then removed under a high vacuum. The crude product was purified by preparative HPLC. Thus, the product having a molecular weight of
20 401.47 (C₂₃H₂₃N₅O₂); MS (ESI): 402 (M+H⁺) was obtained.

1-[4-(2,4-Dimethoxybenzylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea

The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-phenoxyphenyl)urea and 2,4-dimethoxybenzylamine (60°C, 12 h), as described in Example 4. Melting point (ethyl acetate): 214-216°C.

Example 18 1-[1-(2-Dimethylaminoethyl)-2,3-dimethyl-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea



The compound was prepared from 1-(2-dimethylaminoethyl)-2,3-dimethyl-
5 1H-indol-5-ylamine and 4-phenoxyaniline, as described in Example 1. The crude product was purified by preparative HPLC. Thus, the product having a molecular weight of 442.57 ($C_{27}H_{30}N_4O_3$); MS (ESI): 443 ($M+H^+$) was obtained.


1-(2-Dimethylaminoethyl)-2,3-dimethyl-1H-indol-5-ylamine

The compound was obtained by hydrogenation of [2-(2,3-dimethyl-5-
10 nitroindol-1-yl)ethyl]dimethylamine, as described in Example 3. Thus, the product having a molecular weight of 231.34 ($C_{14}H_{21}N_3$); MS (ESI): 232 ($M+H^+$) was obtained.

[2-(2,3-Dimethyl-5-nitroindol-1-yl)ethyl]dimethylamine

Sodium hydride (50% strength in oil; 0.8 g) was added to 2,3-dimethyl-5-
15 nitro-1H-indole (1 g) in tetrahydrofuran (10 mL) at 0°C. After 30 minutes at room temperature, dimethylaminoethyl chloride (hydrochloride; 1.1 g) was added and the mixture was then heated at 65°C for two hours. The cooled reaction solution was extracted with dichloromethane. The organic phase was dried and concentrated. The crude product was purified via chromatography on silica gel
20 (eluent: dichloromethane/methanol 9:1). Thus, the product having a molecular weight of 261.33 ($C_{14}H_{19}N_3O_2$); MS (ESI): 262 ($M+H^+$) was obtained.

Example 19 1-[1-(2-Dimethylaminoethyl)-2-methyl-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea



The compound was prepared from 1-(2-dimethylaminoethyl)-2-methyl-1H-indol-5-ylamine and 4-phenoxyaniline, as described in Example 1. The crude product was purified by preparative HPLC. Thus, the product having a molecular weight of 428.54 ($C_{26}H_{28}N_4O_3$); MS (ESI): 428 ($M+H^+$) was obtained.


1-(2-Dimethylaminoethyl)-2-methyl-1H-indol-5-ylamine

The compound was obtained by hydrogenation of [2-(2-methyl-5-nitroindol-1-yl)ethyl]dimethylamine, as described in Example 3. Thus, the product having a molecular weight of 217.32 ($C_{13}H_{19}N_3$); MS (ESI): 218 ($M+H^+$) was obtained.


[2-(2-Methyl-5-nitroindol-1-yl)ethyl]dimethylamine

The compound was prepared from 2-methyl-5-nitro-1H-indole and dimethylaminoethyl chloride (hydrochloride) as in Example 18. Thus, the product having a molecular weight of 247.30 ($C_{13}H_{17}N_3O_2$); MS (ESI): 248 ($M+H^+$) was obtained.

Table 3: Examples of formula I




where the moiety x_1 is














5







and x_2 is








and x_2 is listed in the column denoted "aniline" of the table below.






Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
20	1-[4-(Cyclohexylmethylamino)phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C26H35N5O	433.60	434
21	1-[4-(Cyclohexylmethylamino)phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C26H35N5O	433.60	434
22	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-pyrrolidin-1-ylphenyl)urea		C23H29N5O	391.52	392
23	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2,5-dimethylpyrrolidin-1-yl)phenyl]urea		C25H33N5O	419.57	420
24	1-[4-(3,6-Dihydro-2H-pyridin-1-yl)phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C24H29N5O	403.53	404
25	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2,6-dimethylmorpholin-4-yl)phenyl]urea		C25H33N5O2	435.57	436







Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
26	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-thiomorpholin-4-yl-phenyl)urea		C23H29N5OS	423.58	424
27	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2-methylpiperidin-1-yl)phenyl]urea		C25H33N5O	419.57	420
28	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2-ethylpiperidin-1-yl)phenyl]urea		C26H35N5O	433.60	434
29	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(3-methylpiperidin-1-yl)phenyl]urea		C25H33N5O	419.57	420
30	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(3,3-dimethylpiperidin-1-yl)phenyl]urea		C26H35N5O	433.60	434

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
31	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(3,5-dimethylpiperidin-1-yl)phenyl]urea		C ₂₆ H ₃₅ N ₅ O	433.60	434
32	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(4-phenylpiperidin-1-yl)phenyl]urea		C ₃₀ H ₃₅ N ₅ O	481.65	482
33	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(4-methylpiperidin-1-yl)phenyl]urea		C ₂₅ H ₃₃ N ₅ O	419.57	420
34	1-(4-Azepan-1-ylphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₅ H ₃₃ N ₅ O	419.57	420
35	1-[4-(Benzylmethylamino)phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₇ H ₃₁ N ₅ O	441.58	442
36	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3[4-(methylphenethylamino)-phenyl]urea		C ₂₈ H ₃₃ N ₅ O	455.61	456

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
37	1-[4-(Butylmethyl- amino)phenyl]-3-[1-(2- dimethylaminoethyl)- 1H-indol-5-yl]urea		C ₂₄ H ₃₃ N ₅ O	407.56	408
38	1-[4-(Benzylbutyl- amino)phenyl]-3-[1-(2- dimethylaminoethyl)- 1H-indol-5-yl]urea		C ₃₀ H ₃₇ N ₅ O	483.66	484
39	1-(4-Dibutylaminophenyl)- 3-[1-(2- dimethylaminoethyl)- 1H-indol-5-yl]urea		C ₂₇ H ₃₉ N ₅ O	449.64	450
40	1-[1-(2-Dimethyl- aminoethyl)-1H-indol- 5-yl]-3-[(4aR,8aS)-4- (octahydroisoquinolin- 2-yl)phenyl]urea		C ₂₈ H ₃₇ N ₅ O	459.64	460
41	1-[1-(2-Dimethylaminoethyl)- 1H-indol-5-yl]-3-[4-(2- methylpyrrolidin-1- yl)phenyl]urea		C ₂₄ H ₃₁ N ₅ O	405.55	406

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
42	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(5-ethyl-2-methylpiperidin-1-yl)phenyl]urea		C27H37N5O	447.63	448
43	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(methylpyridin-3-ylmethylamino)phenyl]urea		C26H30N6O	442.57	443
44	1-[4-(3-Azabicyclo[3.2.2]non-3-yl)phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C27H35N5O	445.61	446
45	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2-isopropylpyrrolidin-1-yl)phenyl]urea		C26H35N5O	433.60	434
46	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2-isobutylpyrrolidin-1-yl)phenyl]urea		C27H37N5O	447.63	448
47	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(3-phenylpyrrolidin-1-yl)phenyl]urea		C29H33N5O	467.62	468

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
48	1-[1-(2-Dimethyl-aminoethyl)-1H-indol-5-yl]-3-[4-(3-trifluoromethylpiperidin-1-yl)-phenyl]urea		C25H30F3N5O	473.55	474
49	1-[1-(2-Dimethyl-aminoethyl)-1H-indol-5-yl]-3-[(4aR,8aR)-4-(octahydroisoquinolin-2-yl)phenyl]urea		C28H37N5O	459.64	460
50	1-[4-(3,4-Dihydro-1H-isoquinolin-2-yl)-phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C28H31N5O	453.59	454
51	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-((1S,5R)-1,3,3-trimethyl-6-azabicyclo[3.2.1]oct-6-yl)phenyl]urea		C29H39N5O	473.67	474
52	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-2,6-dimethylphenyl)urea		C25H34N4O2	422.58	423






Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
53	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-3-methoxyphenyl)urea		C ₂₄ H ₃₂ N ₄ O ₃	424.55	425
54	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-2-methylphenyl)urea		C ₂₄ H ₃₂ N ₄ O ₂	408.55	409
55	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-2,5-dimethylphenyl)urea		C ₂₅ H ₃₄ N ₄ O ₂	422.58	423
56	1-(3,5-Dichloro-4-isobutoxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₃ H ₂₈ Cl ₂ N ₄ O	463.41	463
57	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-3-nitrophenyl)urea		C ₂₃ H ₂₉ N ₅ O ₄	439.52	440
58	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-3-methylphenyl)urea		C ₂₄ H ₃₂ N ₄ O ₂	408.55	409

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
59	1-Benzyl-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]-1-(4-isobutoxyphenyl)urea		C30H36N4O2	484.65	485
60	1-(3-Chloro-4-isobutoxy-5-methylphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C24H31ClN4O2	442.99	443
61	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-2-nitrophenyl)urea		C23H29N5O4	439.52	440
62	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-2,3-dimethylphenyl)urea		C25H34N4O2	422.58	423
63	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(2-fluoro-4-isobutoxyphenyl)urea		C23H29FN4O2	412.51	413






Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
64	1-(3-Chloro-4-isobutoxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C23H29ClN4O2	428.97	429
65	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(3-fluoro-4-isobutoxyphenyl)urea		C23H29FN4O2	412.51	413
66	1-(2-Chloro-4-isobutoxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C23H29ClN4O2	428.97	429
67	Methyl 5-{3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]ureido}-2-isobutoxy benzoate		C25H32N4O4	452.56	453
68	1-(3-Cyano-4-isobutoxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C24H29N5O2	419.53	420
69	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-3,5-dimethylphenyl)urea		C25H34N4O2	422.58	423







Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
70	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-2-trifluoromethylphenyl) urea		C24H29F3N4O	462.52	463
71	1-(4-Butylphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C23H30N4O	378.52	379
72	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxyphenyl)urea		C23H30N4O2	394.52	395
73	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(pyridin-3-yloxy)phenyl]urea		C24H25N5O2	415.50	416
74	1-(3-Cyclopentyloxy-4-methoxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C25H32N4O3	436.56	437

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
75	1-(4-Benzenesulfonyl-2-nitrophenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C25H25N5O5S	507.57	508
76	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2-methoxyphenoxy)-phenyl]urea		C26H28N4O3	444.54	445
77	1-[4-(3-Chlorophenoxy)-phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C25H25ClN4O2	448.96	449
78	1-Biphenyl-4-yl-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C25H26N4O	398.51	399
79	1-[1-(2-Dimethyl-aminoethyl)-1H-indol-5-yl]-3-(2-methoxy-4-phenylaminophenyl)-urea		C26H29N5O2	443.55	444
80	1-(4-Benzyloxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C26H28N4O2	428.54	429



Ex- am- ple	Name	Aniline	Molecular formula	Mol.- weight	[M+H] ⁺
81	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4'-fluorobiphenyl-4-yl)urea		C ₂₅ H ₂₅ FN ₄ O	416.50	417
82	1-(4-Benzylphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₆ H ₂₈ N ₄ O	412.54	413
83	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-pyridin-4-ylmethylphenyl)urea		C ₂₅ H ₂₇ N ₅ O	413.53	414
84	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-p-tolyloxyphenyl)urea		C ₂₆ H ₂₈ N ₄ O ₂	428.54	429
85	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-phenylsulfanylphenyl) urea		C ₂₅ H ₂₆ N ₄ OS	430.58	431

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
86	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(3-trifluoromethyl-phenoxy)phenyl]urea		C26H25F3N4O	482.51	483
87	1-(4-Butyl-2-methylphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C24H32N4O	392.55	393
88	1-(4'-Cyanobiphenyl-4-yl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C26H25N5O	423.52	424
89	1-[4-(4-Chlorophenoxy)-2-trifluoromethylphenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C26H24ClF3N4	516.95	517
90	1-[3-Chloro-4-(pyrimidin-2-yloxy)phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C23H23ClN6O2	450.93	451
91	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(5-methoxy-2-methylbiphenyl-4-yl)urea		C27H30N4O2	442.57	443

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
92	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(piperidine-1-sulfonyl)phenyl]urea		C ₂₄ H ₃₁ N ₅ O ₃ S	469.61	470
93	Ethyl 5-(4-{3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]ureido}phenyl)-2-methylfuran-3-carboxylate		C ₂₇ H ₃₀ N ₄ O ₄	474.56	475
94	1-(4-Benzoxazol-2-ylphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₆ H ₂₅ N ₅ O ₂	439.52	440
95	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(piperidine-1-carbonyl)phenyl]urea		C ₂₅ H ₃₁ N ₅ O ₂	433.56	434
96	1-[3-Cyano-4-(3-trifluoromethylphenyl)sulfanyl]phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₇ H ₂₄ F ₃ N ₅ O	523.58	524

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
97	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-heptafluoropropyl-sulfanylphenyl)urea		C22H21F7N4O	522.49	523
98	1-(4-Benzenesulfonyl-3-chlorophenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C25H25ClN4O3	497.02	497
99	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(pyrimidin-2-yloxy)phenyl]urea		C23H24N6O2	416.49	417
100	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(2-methoxybiphenyl-4-yl)urea		C26H28N4O2	428.54	429
101	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(6-methoxybiphenyl-3-yl)urea		C26H28N4O2	428.54	429
102	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-[1,3]dithiolan-2-ylphenyl)urea		C22H26N4OS2	426.61	427

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
103	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(thiophen-2-ylsulfanyl)phenyl]urea		C ₂₃ H ₂₄ N ₄ OS ₂	436.60	437
104	3-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-1-(4-methoxyphenyl)-1-methylurea		C ₂₁ H ₂₆ N ₄ O ₂	366.47	367
105	1-[4-(2-Chlorophenoxy)-phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₅ H ₂₅ ClN ₄ O ₂	448.96	449
106	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(6-phenoxy-pyridin-3-yl)urea		C ₂₄ H ₂₅ N ₅ O ₂	415.50	416
107	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-m-tolyloxyphenyl)urea		C ₂₆ H ₂₈ N ₄ O ₂	428.54	429

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
108	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-o-tolyloxyphenyl)urea		C ₂₆ H ₂₈ N ₄ O ₂	428.54	429
109	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(3-methoxyphenoxy)-phenyl]urea		C ₂₆ H ₂₈ N ₄ O ₃	444.54	445

The molecule ion peak $[(M+H)^+]$ was taken from ESI mass spectra.

The examples 20 – 51 and 71 -109 were prepared according to Example 1.

5 Synthesis of Examples 52 – 70

Carbonyldiimidazole (0.25 mmol) was added to 1-(2-dimethylaminoethyl)-1H-indol-5-ylamine (0.25 mmol) in dimethylformamide (1 mL) at 0°C. After 1 hour at room temperature, the reaction solution was cooled again to 0°C and the appropriate aminophenol (0.25 mmol) was added. After 15 hours at room temperature, cesium carbonate (0.5 mmol) and isobutyl iodide (0.5 mmol) were added and the solution was heated at 80°C for 2 hours. The reaction solutions were filtered and the filtrate was washed with sodium bicarbonate (5% strength) and sodium chloride solution (5% strength). The organic phase was dried and concentrated. The crude product was purified by preparative HPLC. Thus, the product having the molecular weight indicated in Table 3 and the molecule ion peak of the mass spectrum, likewise indicated in Table 3, was obtained.

Precursors of Examples 20-51

A mixture of 4-fluoronitrobenzene (0.35 mmol), potassium carbonate (0.7 mmol), the appropriate amine and dimethylformamide (1 mL) was heated to 100°C for three hours. The reaction solution was filtered and washed with sodium chloride solution (5% strength). The organic phase was dried and concentrated. The 4-nitroaniline obtained as crude product was dissolved in glacial acetic acid (1 mL) and zinc dust (0.25 g) was added. After a reaction time of 3 hours, the reaction solution was diluted with ethyl acetate (10 mL), filtered and the filtrate was washed with sodium chloride solution (5% strength). The filtrate was dried and concentrated. The obtained crude product, 4-substituted aniline, was reacted further without any further purification.

The following 4-nitroanilines were prepared:

- 1-(4-nitrophenyl)azocan
- cyclohexylmethyl-(4-nitrophenyl)amine
- 15 1-(4-nitrophenyl)pyrrolidine
- 2,5-dimethyl-1-(4-nitrophenyl)pyrrolidine
- 1-(4-nitrophenyl)-1,2,3,6-tetrahydropyridine
- 2,6-dimethyl-4-(4-nitrophenyl)morpholine
- 4-(4-nitrophenyl)thiomorpholine
- 20 2-methyl-1-(4-nitrophenyl)piperidine
- 2-ethyl-1-(4-nitrophenyl)piperidine
- 3-methyl-1-(4-nitrophenyl)piperidine
- 3,3-dimethyl-1-(4-nitrophenyl)piperidine
- 3,5-dimethyl-1-(4-nitrophenyl)piperidine

- 1-(4-nitrophenyl)-4-phenylpiperidine
- 4-methyl-1-(4-nitrophenyl)piperidine
- 2-(4-nitrophenyl)-1,2,3,4-tetrahydroisoquinoline
- 1-(4-nitrophenyl)azepan
- 5 benzylmethyl-(4-nitrophenyl)amine
- methyl-(4-nitrophenyl)phenethylamine
- butylmethyl-(4-nitrophenyl)amine
- benzylbutyl-(4-nitrophenyl)amine
- dibutyl-(4-nitrophenyl)amine
- 10 (4aR,8aS)-2-(4-nitrophenyl)decahydroisoquinoline
- 2-methyl-1-(4-nitrophenyl)pyrrolidine
- 5-ethyl-2-methyl-1-(4-nitrophenyl)piperidine
- methyl-(4-nitrophenyl)pyridine-3-ylmethylamine
- 3-(4-nitrophenyl)-3-azabicyclo[3.2.2]nonane
- 15 2-isopropyl-1-(4-nitrophenyl)pyrrolidine
- 2-isobutyl-1-(4-nitrophenyl)pyrrolidine
- 1-(4-nitrophenyl)-3-phenylpyrrolidine
- 1-(4-nitrophenyl)-3-trifluoromethylpiperidine
- (4aR,8aR)-2-(4-nitrophenyl)decahydroisoquinoline
- 20 (1S,5R)-1,3,3-trimethyl-6-(4-nitrophenyl)-6-azabicyclo[3.2.1]octane

All of the 4-nitroanilines listed above showed the expected molecule ion peak in the ESI mass spectrum.

The following 4-substituted anilines were prepared:

4-azocan-1-ylphenylamine

N-cyclohexyl-N-methylbenzene-1,4-diamine

4-pyrrolidin-1-ylphenylamine

5 4-(2,5-dimethylpyrrolidin-1-yl)phenylamine

4-(3,6-dihydro-2H-pyridin-1-yl)phenylamine

4-(2,6-dimethylmorpholin-4-yl)phenylamine

4-thiomorpholin-4-ylphenylamine

4-(2-methylpiperidin-1-yl)phenylamine

10 4-(2-ethylpiperidin-1-yl)phenylamine

4-(3-methylpiperidin-1-yl)phenylamine

4-(3,3-dimethylpiperidin-1-yl)phenylamine

4-(3,5-dimethylpiperidin-1-yl)phenylamine

4-(4-phenylpiperidin-1-yl)phenylamine

15 4-(4-methylpiperidin-1-yl)phenylamine

4-(3,4-dihydro-1H-isoquinolin-2-yl)phenylamine

4-azepan-1-ylphenylamine

N-benzyl-N-methylbenzene-1,4-diamine

N-methyl-N-phenethylbenzene-1,4-diamine

20 N-butyl-N-methylbenzene-1,4-diamine

N-benzyl-N-butylbenzene-1,4-diamine

N,N-dibutylbenzene-1,4-diamine

(4aR,8aS)-4-(octahydroisoquinolin-2-yl)phenylamine

4-(2-methylpyrrolidin-1-yl)phenylamine

4-(5-ethyl-2-methylpiperidin-1-yl)phenylamine

5 N-methyl-N-pyridin-3-ylmethylbenzene-1,4-diamine

4-((1S,5R)-1,3,3-trimethyl-6-azabicyclo[3.2.1]oct-6-yl)phenylamine

4-(3-azabicyclo[3.2.2]non-3-yl)phenylamine

4-(2-isopropylpyrrolidin-1-yl)phenylamine

4-(2-isobutylpyrrolidin-1-yl)phenylamine


10 4-(3-phenylpyrrolidin-1-yl)phenylamine

4-(3-trifluoromethylpiperidin-1-yl)phenylamine

(4aR,8aR)-4-(octahydroisoquinolin-2-yl)phenylamine.

All of the 4-substituted anilines listed above showed the expected molecule ion peak in the ESI mass spectrum.


15 **Example 110 4-Phenoxyphenyl [1-(2-dimethylaminoethyl)-1H-indol-5-yl]carbamate**



The compound was prepared according to Example 1 by reacting the carbonyldiimidazole-activated indolamine with deprotonated 4-phenoxyphenol.

20 Thus, the product having a molecular weight of 415.50 (C₂₅H₂₅N₃O₃); MS (ESI): 416 (M+H⁺) was obtained.

Example 111 1-(2-Imidazol-1-ylmethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea



5 Mesyl chloride (47 μ l) was added to 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea (0.2 g) and triethylamine (0.16 mL) in dichloromethane (4 mL) at 0°C. After 10 minutes, imidazole (185 mg) was added. After 12 hours, the reaction solution was washed with sodium chloride solution, dried and concentrated. The crude product was purified by preparative HPLC.
10 Thus, the product having a molecular weight of 437.51 ($C_{26}H_{23}N_5O_2$); MS (ESI): 438 ($M+H^+$) was obtained.


1-(2-Hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea

(5-Amino-1-methyl-1H-indol-2-yl)methanol was reacted with 4-phenoxyaniline and carbonyldiimidazole, as described in Example 1. Thus, the
15 product having a molecular weight of 387.44 ($C_{23}H_{21}N_3O_3$); MS (ESI): 388 ($M+H^+$) was obtained.

(5-Amino-1-methyl-1H-indol-2-yl)methanol


(1-Methyl-5-nitro-1H-indol-2-yl)methanol was hydrogenated as described in Example 3. Thus the product having a molecular weight of 176.22 ($C_{10}H_{12}N_2O$);
20 MS (ESI): 177 ($M+H^+$) was obtained.

Example 112 1-[1-Methyl-2-(2-methyl-4,5-dihydroimidazol-1-ylmethyl)-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea



The compound was prepared from 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea and 2-methyl-4,5-dihydroimidazole, as described in Example 111. Thus, the product having a molecular weight of 453,55 ($C_{27}H_{27}N_5O_2$); MS (ESI): 454 ($M+H^+$) was obtained.


Example 113 1-(2-Cyclohexylaminomethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea



10

The compound was prepared from 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea and cyclohexylamine, as described in Example 111. Thus, the product having a molecular weight of 468.60 ($C_{29}H_{32}N_4O_2$); MS (ESI): 469 ($M+H^+$) was obtained.


Example 114 1-[2-(3-Dimethylaminopyrrolidin-1-ylmethyl)-1-methyl-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea



The compound was prepared from 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea and 3-dimethylaminopyrrolidine, as described in Example 111. Thus, the product having a molecular weight of 483.62 ($C_{29}H_{33}N_5O_2$); MS (ESI): 484 ($M+H^+$) was obtained.


5

Example 115 1-[2-(4-Hydroxypiperidin-1-ylmethyl)-1-methyl-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea



The compound was prepared from 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea and 4-hydroxypiperidin, as described in Example 111. Thus, the product having a molecular weight of 470.58 ($C_{28}H_{30}N_4O_3$); MS (ESI): 471 ($M+H^+$) was obtained.


Example 116 1-[1-Methyl-2-(4-phenylpiperidin-1-ylmethyl)-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea



15


The compound was prepared from 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea and 4-phenylpiperidine, as described in Example 111. Thus, the product having a molecular weight of 530.68 ($C_{34}H_{34}N_4O_2$); MS (ESI): 531 ($M+H^+$) was obtained.

Example 117 N-(1-{1-Methyl-5-[3-(4-phenoxyphenyl)ureido]-1H-indol-2-ylmethyl}pyrrolidin-3-yl)acetamide



The compound was prepared from 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea and pyrrolidin-3-ylacetamide, as described in
5 Example 111. Thus, the product having a molecular weight of 497.60 ($C_{29}H_{31}N_5O_3$); MS (ESI): 498 ($M+H^+$) was obtained.

Example 118 1-(4-Phenoxyphenyl)-3-(2-pyrrolidin-1-ylmethylbenzofuran-5-yl)urea



10

The compound was prepared from 2-pyrrolidin-1-ylmethylbenzofuran-5-ylamine and 4-phenoxyaniline, as described in Example 1. Thus, the product having a molecular weight of 427.51 ($C_{26}H_{25}N_3O_3$); MS (ESI): 428 ($M+H^+$) was obtained.

15 2-Pyrrolidin-1-ylmethylbenzofuran-5-ylamine

The compound was prepared by hydrogenation of 1-(5-nitrobenzofuran-2-ylmethyl)pyrrolidine, as described in Example 3. Thus, the product having a molecular weight of 216.29 ($C_{13}H_{16}N_2O$); MS (ESI): 217 ($M+H^+$) was obtained.

20


1-(5-Nitrobenzofuran-2-ylmethyl)pyrrolidine

The compound was prepared from (5-nitrobenzofuran-2-yl)methanol, as described in Example 3. Thus, the product having a molecular weight of 246.27 ($C_{13}H_{14}N_2O_3$); MS (ESI): 247 ($M+H^+$) was obtained.

5 (5-Nitrobenzofuran-2-yl)methanol

The compound was prepared by reduction of methyl 5-nitrobenzofuran 2-carboxylate, as described in Example 3. Thus, the product having a molecular weight of 193.16 ($C_9H_7NO_4$); MS (ESI): 194 ($M+H^+$) was obtained.

10 **Example 119 1-(4-Phenoxyphenyl)-3-(2-pyrrolidin-1-ylmethylbenzo[b]thiophen-5-yl)urea**



15 The compound was prepared from 2-pyrrolidin-1-ylmethylbenzo[b]thiophen-5-ylamine and 4-phenoxyaniline, as described in Example 1. Thus, the product having a molecular weight of 443.57 ($C_{26}H_{25}N_3O_2S$); MS (ESI): 444 ($M+H^+$) was obtained.

2-Pyrrolidin-1-ylmethylbenzo[b]thiophen-5-ylamine

20 The compound was prepared by hydrogenation of 1-(5-nitrobenzo[b]thiophen-2-ylmethyl)pyrrolidine, as described in Example 3. Thus, the product having a molecular weight of 232.35 ($C_{13}H_{16}N_2S$); MS (ESI): 233 ($M+H^+$) was obtained.

1-(5-Nitrobenzo[b]thiophen-2-ylmethyl)pyrrolidine

The compound was prepared from (5-nitrobenzo[b]thiophen-2-yl)methanol, as described in Example 3. Thus, the product having a molecular weight of 262.33 ($C_{13}H_{14}N_2O_2S$); MS (ESI): 263 ($M+H^+$) was obtained.

(5-Nitrobenzo[b]thiophen-2-yl)methanol

The compound was prepared by reduction of methyl 5-nitrobenzo[b]thiophene 2-carboxylate, as described in Example 3. Thus, the product having a molecular weight of 209.23 ($C_9H_7NO_3S$); MS (ESI): 210 ($M+H^+$)
5 was obtained.

In general, all of the basic compounds described were obtained either as free bases or in the form of a salt of one of the following acids: formic acid, trifluoroacetic acid or hydrochloric acid.